

# “BLISTER BEETLE JUICE” WOUNDCARE INSTRUCTIONS

- Avoid getting the treated area moist with sweat or water until after cantherone has been washed off.
- Wash the cantherone off in \_\_\_\_\_ hours. If a bandage was applied, soak it in water before removing to ease removal. Do not scrub but ensure all cantherone is removed to prevent further blistering. Apply Vaseline or Polysporin (NOT NEOSPORIN) to the site and cover with a bandage twice daily. Some patients develop an irritant or allergic reaction to bandages. If this occurs, use paper tape and gauze.
- It is normal for the site to be a red and develop a blister. You may pop the blister with a sterile needle and drain the fluid if it becomes uncomfortable. If you notice any signs of infection such as fever, foul smelling drainage, or marked redness, swelling, or intense pain, call our office.
- DO NOT ALLOW THE CANTHERONE TO SPREAD OR RUB ONTO OTHER AREAS OF YOUR BODY AS EVERY PLACE THAT COMES INTO CONTACT WITH THE CANTHERONE WILL BLISTER. THIS CAN OCCUR WHEN YOU SWEAT OR AN UNTREATED AREA OF SKIN TOUCHES A TREATED AREA. BE ESPECIALLY CAREFUL IF THE TREATMENT AREA IS IN A CREASE OF YOUR SKIN (i.e. BACK OF KNEES)
- This method will require several treatments to fully resolve.